

Homoeostasis in Neurotoxicity. The Case of Organotin Salts

· E-mail Article

M. Marinovich^a, B. Viviani^a and C. L. Galli^a

Available online 14 January 1998.

Abstract

The cytoskeleton is a major constituent of the neuronal cytoplasm; it controls cell shape and plays important roles in regulating various physiological processes. In neurons, actin filaments are involved in the growth of the neurite and the neurotransmitter release. Recent findings suggest that actin filaments play a role in modulating [Ca²⁺]; responses to neurotoxic insults. The physiological functioning of the neural cell is critically dependent on the intracellular distribution of calcium. An increase of cytosolic free calcium can activate a number of intracellular reactions, including neurotransmitter release, protein phosphorylation, protease activity, and, eventually, cell death. Many neurotoxic agents with diverse mechanisms have been reported to affect mechanisms associated with calcium. Among these are organotin compounds: they can both raise the cytosolic and synaptosomal [Ca²⁺]; concentrations and interfere with the [Ca²⁺]; response evoked by different agonists. Furthermore, some of these compounds cause actin depolymerization. The interference of triethyltin (TET)--a compound inducing myelin vacuolization and brain oedema--with Ca²⁺ homoeostasis and actin polymerization results in an adverse effect on neurotransmitter release in different neural cell lines. However, another neurotoxic organotin compound (trimethyltin, TMT) induces apoptosis in neural cells through the activation of a Ca²⁺ dependent pathway. In conclusion, the identification of the key changes in actin and Ca²⁺ homoeostasis could give early information on neural cell perturbation resulting in altered functionality or even cell death.

^a Laboratory of Toxicology, Institute of Pharmacological Sciences, University of Milan, via Balzaretti 9 20133 Milan Italy

HOME HELP FEEDBACK SUBSCRIPTIONS ARCHIVE SEARCH TABLE OF CONTENTS

| SCIENCE DIRECT Register or Login: US | er name Password: Go , |
|---|--|
| Home Journals Abstract Databases Reference Wo | My Alerts My Profile Help WELCOME GUEST USER [Info] |
| return to SCIfUS Survey of Ophthalmology Volume 45, Supplement 3, May 2001, Pages S250-S255 | |
| doi:10.1016/S0039-6257(01)00198-9 | This Document Abstract Full Text + Links PDF (65 K) |
| Original article Now Howizons in Novement action | Actions E-mail Article |

New Horizons in Neuroprotection

Rita Naskar PhD and Evan B. Dreyer MD, PhD

Scheie Eye Institute and Department of Ophthalmology, the University of Pennsylvania, Philadelphia, PA, USA

Available online 2 November 2001.

Abstract

Glaucoma is a leading cause of blindness worldwide and the second leading cause of irreversible blindness in the USA. The most common form of glaucoma, primary open angle glaucoma, is characterized by a chronically elevated intraocular pressure in the absence of any demonstrable structural abnormalities in the eye. The pathologic hallmark of glaucomatous optic neuropathy is the selective death of retinal ganglion cells associated with structural changes in the optic nerve head. Recent discoveries suggest a role for nitric oxide, glutamate, apoptosis, and others, in the pathophysiology of this neuropathy. These newer discoveries are addressed in this article.

Author Keywords: apoptosis; caspase; glaucoma; glutamate; nitric oxide; retinal; ganglion cells

Reprint address: Evan Benjamin Dreyer, MD, PhD, Scheie Eye Institute, 51 North 39th Street, Philadelphia, PA 19104 USA

Survey of Ophthalmology

Volume 45, Supplement 3, May 2001, Pages S250-S255

This Document

- **▶** Abstract
- Full Text + Links
- · PDF (65 K)